

Journal of Fish and Wildlife Management

Patterns of Florida Bonneted Bat Occupancy at the Northern Extent of Its Range

--Manuscript Draft--

Manuscript Number:	JFWM-22-055R1
Article Type:	Article
Keywords:	Eumops floridanus; Florida bonneted bat; forest cover; longleaf pine; red-cockaded woodpecker
Corresponding Author:	Robert A. Schorr Colorado State University, CNHP Fort Collins, Colorado UNITED STATES
First Author:	Robert A. Schorr
Order of Authors:	Robert A. Schorr Kristopher A. Pitcher Robert A. Aldredge Paul M. Lukacs
Abstract:	<p>The Florida bonneted bat <i>Eumops floridanus</i> is a rare, endemic bat of south Florida, which roosts in woodpecker cavities, and anthropogenic structures, such as roofing tiles, chimneys, and bat houses. The northern-most occurrences of the bonneted bat are from mature pine forests at the Avon Park Air Force Range, Florida. We used ultrasonic acoustic recorders to understand bonneted bat activity and habitat occupancy. We modeled occupancy using a hierarchical Bayesian analysis, and included site- and time-specific covariates of detection probability, and site-specific covariates of occupancy. Probability of detection was low throughout Avon Park Air Force Range, but increased with Julian date. In most habitats, occupancy was poorly estimated, except for flatwood mature pinelands where occupancy was low (0.23 ± 0.06). As distance from red-cockaded woodpecker colonies increased occupancy decreased ($\beta = -1.19 \pm 0.26$ SD). At the northern-most extent of the range, and throughout much of the historic range, increasing the expanse of mature, fire-maintained forest systems will increase habitat for the bonneted bat, and lead to faster population recovery.</p>

Running Heading: Florida bonneted bat occupancy

1 **Patterns of Florida Bonneted Bat Occupancy at the Northern Extent of Its Range**

2

3 Robert A. Schorr,* Kristopher A. Pitcher, Robert A. Aldredge, Paul M. Lukacs

4

5 **R.A. Schorr**

6 Colorado Natural Heritage Program, Colorado State University, Fort Collins, CO 80523

7 **K.A. Pitcher, R.A. Aldredge**

8 U.S. Fish and Wildlife Service, Avon Park Air Force Range, Avon Park, FL 33825

9 **P.M. Lukacs**

10 Wildlife Biology Program, W.A. Franke College of Forestry and Conservation, University of

11 Montana, Missoula, MT 59812

12

13 *Corresponding author: robert.schorr@colostate.edu

14

15

Abstract

16
17 The Florida bonneted bat *Eumops floridanus* is a rare, endemic bat of south Florida, which roosts
18 in woodpecker cavities, and anthropogenic structures, such as roofing tiles, chimneys, and bat
19 houses. The northern-most occurrences of the bonneted bat are from mature pine forests at the
20 Avon Park Air Force Range, Florida. We used ultrasonic acoustic recorders to understand
21 bonneted bat activity and habitat occupancy. We modeled occupancy using a hierarchical
22 Bayesian analysis, and included site- and time-specific covariates of detection probability, and
23 site-specific covariates of occupancy. Probability of detection was low throughout Avon Park
24 Air Force Range, but increased with Julian date. In most habitats, occupancy was poorly
25 estimated, except for flatwood mature pinelands where occupancy was low (0.23 ± 0.06). As
26 distance from red-cockaded woodpecker colonies increased occupancy decreased ($\beta = -1.19 \pm$
27 0.26 SD). At the northern-most extent of the range, and throughout much of the historic range,
28 increasing the expanse of mature, fire-maintained forest systems will increase habitat for the
29 bonneted bat, and lead to faster population recovery.

30
31 Key words: *Eumops floridanus*, Florida bonneted bat, forest cover, longleaf pine, red-cockaded
32 woodpecker

33
34 Received: September 2022; Accepted: May 2023; Published Online Early: July 2023; Published:
35 December 2023

36

37 Citation: Schorr RA, Pitcher KA, Aldredge RA, Lukacs PM. 2023. Patterns of Florida bonneted
38 bat occupancy at the northern extent of its range. *Journal of Fish and Wildlife Management*
39 14(2):xx-xx; e1944-687X. <https://doi.org/10.3996/JFWM-22-055>

40
41 This Online Early paper will appear in its final typeset version in a future issue of the *Journal of*
42 *Fish and Wildlife Management*. This article has been accepted for publication and undergone full
43 peer review but has not been through the copyediting, typesetting, pagination, and proofreading
44 process, which may lead to differences between this version and the Version of Record. The
45 findings and conclusions in this article are those of the author(s) and do not necessarily represent
46 the views of the U.S. Fish and Wildlife Service.

47

48 Introduction

49 The Florida bonneted bat *Eumops floridanus* (herein called “bonneted bat”) is a federally
50 endangered species endemic to south Florida, with a limited range in seven counties (US
51 Endangered Species Act (ESA 1973, as amended); USFWS 2013). It is a large (40 – 65 g; Figure
52 1) bat that is active year-round and roosts in anthropogenic structures, such as tile roofing,
53 chimney structures, or bat houses, and in palms, and woodpecker cavities of live mature pines
54 and dead pine snags (Angell and Thompson 2015; Gore et al. 2015; Braun de Torrez et al. 2016;
55 Webb et al. 2021). The bonneted bat roosts in small colonies, which makes studying its
56 population ecology and habitat needs challenging (Gore et al. 2015; Bailey et al. 2017a).
57 Bonneted bats will roost in red-cockaded woodpecker *Picoides borealis* (RCW) cavities, and
58 likely used these more frequently when distribution of mature southern pines, such as longleaf
59 pine *Pinus palustris* and slash pine *Pinus elliottii* was greater (Walker 2000). These fire-adapted

60 ecosystems provided ideal open forests for bonneted bats to forage and roost (Angell and
61 Thompson 2015; Braun de Torrez 2018a; Braun de Torrez et al. 2018b).

62 The northern-most roosts of the bonneted bat occur at Avon Park Air Force Range (APAFR),
63 where there are stands of mature longleaf forest (Angell and Thompson 2015). Five active and
64 inactive bonneted bat roosts at APAFR are in RCW cavities. As with RCWs, bonneted bats may
65 have been more prevalent at APAFR when mature longleaf and slash pine forests were
66 ubiquitous in south-central Florida (Belwood 1992). The bonneted bat roosts at APAFR are
67 considerable distances from the nearest bonneted bat population along the Peace River
68 approximately 40 km southeast. The isolation of populations provides representation, and
69 conservation of these populations is critical for recovering bonneted bat populations and delisting
70 the species (Smith et al. 2018; Austin et al. 2022). The discovery of bonneted bats at APAFR
71 (Angell and Thompson 2015) gives promise that other roosts exist in similar forested systems in
72 south-central Florida, and it heightens the importance of conserving each isolated, small
73 population.

74 Current land management at APAFR is a balance between maintaining historic habitat types,
75 managing pine plantations, conserving native species, reducing natural fuel hazards, and
76 conducting daily military activities. The APAFR actively conserves habitat for rare species, like
77 the RCW, Florida scrub jay *Aphelocoma coerulescens*, and Florida grasshopper sparrow
78 *Ammodramus savannarum*, and bonneted bats. To accommodate the varied land use demands,
79 APAFR needs to understand where these rare species are, how they use the available habitats,
80 and how to minimize land management activities that threaten the species and their habitats. For
81 bonneted bats, biologists identify habitat use by collecting ultrasonic acoustic records throughout
82 the range. In this study, we use those recordings to estimate bonneted bat habitat use, and use

83 environmental and location-specific covariates to inform habitat occupancy and detection
84 probability.

85

86 **Study site**

87 We conducted this study at APAFR that is a 42,900-ha air-to-ground training complex
88 with 40,000 ha undeveloped lands in Polk and Highland counties, Florida (elevation = 37 m)
89 (Figure 2). Wildfires regularly occur at APAFR, ignited by lightning strikes and unintentionally
90 by on-going military training. In addition, prescribed fire is used to reduce fuel loads and
91 maintain pyrogenic-adapted land cover types, such as longleaf pine forest, oak *Quercus* scrub,
92 and dry prairie. The application of prescribed fire at APAFR contrasts the surrounding lands
93 where fire was suppressed in past decades. The wet season of APAFR typically lasts from mid-
94 May to November with warm temperatures (18-32°C) and ample precipitation (89 ± 27 cm/yr),
95 and the dry season (November to mid-May) has comparatively cooler temperatures and less
96 precipitation (12-25°C, 42 ± 15 cm/yr; Duever et al. 1994; Slocum et al. 2010).

97 Roosting areas for the bonneted bat at APAFR are a mosaic of hydric flatwoods of south
98 Florida slash pine *P. e. densa* and pine flatwood-savannahs of longleaf pine (Figure 1). The
99 understory vegetation of hydric flatwoods is dominated by cutthroat grass *Panicum abscissum*,
100 while longleaf pine flatwoods and savannahs are dominated by wiregrass *Aristida beyrichiana*
101 and patchy woody shrubs, such as saw palmetto *Serenoa repens*. Other abundant land cover of
102 APAFR includes managed mixed pine plantations of north Florida slash pine, south Florida
103 slash, and long leaf pine, swamps of pond cypress *Taxodium ascendens* and bald cypress *T.*
104 *distichum*, hardwood hammocks, oak and pine scrublands, marshes, and grassland prairies

105 (Figure 2). Less than 1% of APAFR is developed lands that include a cantonment area, roads,
106 and a runway.

107

108 **Methods**

109 **Acoustic call collection and analysis**

110 We conducted acoustic surveys from November 21, 2018, to March 10, 2020, deploying
111 autonomous recording units (ARU) with ultrasonic microphones (SM4BAT acoustic recorders
112 and SMM_U2 microphones; Wildlife Acoustics, Inc., Maynard, MA) at 508 sites. We focused
113 our surveys in areas that had high potential for bonneted bat roosting habitat and areas with
114 higher potential for tree cavities or snags. We surveyed hydric flatwoods, pine flatwoods and
115 savannahs, pine plantations over 15 years old, cypress swamps, hardwood hammocks, and oak
116 and pine scrub, with some random points occurring in nearby grasslands, which encompassed
117 21,000 ha. We randomly selected ARU deployment locations that were separated by >100 m.
118 We mounted ARUs on a pole and tripod, with microphones at the top of the pole (4.9 m) and
119 started recording 30 minutes before sunset and ended 30 minutes after sunrise. We set ARUs to
120 record with a minimum trigger frequency of 8 kHz because bonneted bats emit low frequency
121 calls between 10 to 18 kHz (Belwood 1992). Also, we set ARUs at a sample rate of 256 kHz,
122 volume trigger level at 12dB, minimum trigger window to 2 seconds, and minimum call duration
123 of 2 msec (Braun de Torrez et al. 2018a). We deployed each ARU for >3 consecutive nights.

124 To analyze recorded call files we used Kaleidoscope Pro 5.4.1 (Wildlife Acoustics, Inc.)
125 and the Bats of North America 5.4.0 classifier for Florida (Agranat 2013). We processed call
126 files using the Balanced or Neutral setting to filter out noise, low-quality calls, or non-bat call
127 files (Reichert et al. 2018), then used the auto-identification function software to identify

128 bonneted bats. We defined a call file as a recording that had at least three distinct pulses (bat-
129 emitted ultrasound) lasting between 2 and 15 seconds. We manually vetted all call files identified
130 as bonneted bats, removing any false positives and removing bat social calls, insect noise, and
131 bird vocalizations. We classified a call as coming from bonneted bat when the call had a
132 characteristic frequency of 10-18 kHz, a maximum frequency of 16-22 kHz, average call
133 duration of 10.2ms, and an average call per second of 5.5.

134

135 **Temporal and habitat-specific analysis covariates**

136 We collected regional precipitation and temperature covariates from a weather station on
137 APAFR, and collected site-specific temperature using the summary files generated by the ARU
138 at each survey location. We calculated habitat covariates for each survey location in ArcGIS Pro
139 2.6.5 (ESRI, Redlands, California) using land cover feature layers, and used remote-sensed Light
140 Detection and Ranging (LIDAR) data (October 2018) in R (v 3.5.0) to determine tree stand
141 covariates within 100 m of each ARU (Beucher and Meyer 1993; Popescu and Wynne 2004; R
142 Core Team 2018; Plowright 2020). We identified individual trees and their canopy in the “Forest
143 Tools” package by setting the minimum tree height to 2 m and defining the dynamic window
144 size ($\text{lin} \leftarrow \text{function}(x)\{x*0.05+2\}$; Popescu and Wynne 2004). Using burn history data, we
145 determined the date since last burn at each ARU location.

146 For modeling detection probability (p), we used weather covariates of maximum,
147 minimum, and mean overnight temperature range on ARU environmental data and from a nearby
148 weather station (Avon Park, FL; 27.61°N 81.51°W). We used weather station data of total
149 nightly precipitation, and maximum, minimum, and mean wind speed. We used ARU-specific
150 covariates of total number of call files auto-identified as Brazilian free-tailed bat *Tadarida*

151 *brasiliensis* (TABR) calls and noise. Also, we used habitat covariates that aggregated the major
152 habitat types at APAFR, including mature flatwood pinelands (flatwoods, which included hydric
153 flatwoods and pine flatwoods), pine plantations (plantations), scrubby flatwoods and sandhills
154 (scrub), oak hammock (oak), and cypress swamplands (swamp). An additional habitat type
155 (prairies) was not included in the original sampling design; however, some ARUs were placed in
156 pine plantations that had been clearcut and resembled grassland prairies. We used Julian date
157 because bonneted bats have increased detection probability later in the year, and we created a
158 covariate for mid-winter and mid-summer months when dominant male bonneted bats show
159 higher activity (January – March, July - August; Braun de Torrez et al. 2020).

160 For modeling occupancy, we used covariates of the six major habitat types used to model
161 p, and distances to and sizes of particular land features (Table 1). We used distance to nearest
162 RCW cluster, distance to nearest orange *Citrus sinensis* grove, area of nearest orange grove,
163 distance to nearest wetland, and area of nearest wetland. Also, we used ARU specific landscape
164 measurements within 100-km radius pertaining to forest canopy cover, forest crown area, tree
165 heights, and canopy radius (Table 1). In addition, we used days since last fire as a covariate. To
166 test if bonneted bat detections are lower during periods of high TABR activity, we calculated the
167 number of call files auto-identified as TABR at each ARU and used those as ARU-specific
168 covariates of detection. To detect if high levels of insect noise interfered with our ability to
169 detect bonneted bats, we calculated the number of noise call files recorded at each ARU location.

170

171 **Models and Analysis**

172 We used Bayesian hierarchical occupancy models to estimate occupancy probability
173 assuming imperfect detection (Royle and Darazio 2008; Bailey et al. 2017). We mean-centered

174 and standardized occupancy and detection covariates to speed Markov chain Monte Carlo
175 (MCMC) convergence and modeled detection probability first, while holding occupancy time
176 dependent (Morin et al. 2020). We used JAGS v 4.3.0 launched from RStudio v 1.3.1073 with
177 the R2jags library (Su and Yajima 2021) for Bayesian estimation of model parameters via
178 MCMC samples of posterior distributions. We input each covariate as a random effect using a
179 vague, normally distributed $[N(0,0.01)]$ prior on logit-scale parameters (Kery and Royle 2015).
180 Posterior samples were ranged on 50,000 MCMC samples of posterior distributions of three
181 chains, following a burn-in of 10,000. We assessed convergence of MCMC chains using trace
182 plots and Gelman-Rubin diagnostics (\hat{R}). Convergence was reached for all parameters according
183 to the criteria $|\hat{R} - 1| < 0.1$ (Ntzoufras 2009). We standardized all covariates to speed MCMC
184 convergence.

185 We evaluated models using an indicator variable selection process (Hobbs and Hooten
186 2015). We built parameter weights for covariates, which depict the percentage of time a
187 particular covariate was included in the model iteration (Kuo and Mallick 1998). We ran models
188 with the full suite of covariates and removed any covariates that were included in less than 50%
189 of the iterations. We reran the analysis with covariates that were included in $\geq 50\%$ of iterations.

190

191 Results

192 We conducted acoustic recording at 498 sites at APAFR and deployed ARUs from 3 – 25 nights
193 (mean: 5 nights). Acoustic analysis was not conducted at nine ARUs because of recording errors.
194 Some ARUs recorded for multiple weeks because weather and military activity prevented
195 retrieval. We recorded bonneted bats at 128 locations, and estimated that bonneted bats could be
196 at 166 locations (95% credible interval (CI): 149 – 188). The best model for detection probability

197 included covariates of Julian date and habitat type. Detection probability was comparably low in
198 all habitat types ($p = 0.07 - 0.10, \pm 0.01 - 0.04$ SD), but detection probability was substantially
199 lower in oak habitats ($p = 0.02 \pm 0.01$ SD, Table 2). Including the covariate Julian date increased
200 p ($\beta = 0.41 \pm 0.07$ SD).

201 The best model for occupancy included three habitat types, including pine plantations,
202 prairies, and flatwoods. Occupancy was greatest in plantations ($\psi = 0.39 \pm 0.33$ SD), then
203 prairies ($\psi = 0.38 \pm 0.32$ SD), and flatwoods ($\psi = 0.23 \pm 0.06$ SD), but was poorly estimated in
204 all but flatwood habitats ($CV > 50\%$). Occupancy decreased as distance to a RCW colony
205 increased ($\beta = -1.19 \pm 0.26$ SD, Figure 3). Also, occupancy decreased as the area of nearest
206 wetland increased ($\beta = -0.26 \pm 4.01$ SD), and as distance to nearest orange grove increased ($\beta = -$
207 0.42 ± 3.05 SD), but these impacts are poorly estimated.

208

209

Discussion

210 Bonneted bats were detected in all the APAFR habitat types included in this analysis, including
211 scrub, swamp, flatwoods, prairies, and pine plantations, but only occurred in pine plantations,
212 flatwoods, and prairies often enough to produce estimates of occupancy. Additionally, only
213 occupancy estimates in flatwoods habitats produced reasonable estimates of variability.

214 Flatwoods habitats include all native, old-growth longleaf and slash pine where bonneted bat
215 roosts occur at APAFR. It is these mature pine stands where RCWs and the cavities they build
216 are more available as bonneted bat roosts. Thus, it is not surprising that two of the best predictors
217 of bonneted bat occupancy are the presence of flatwoods habitat types and the distance to RCW
218 clusters. At APAFR there are 45 active clusters and 8 inactive clusters, with 229 artificial
219 cavities and 119 natural cavities in 348 cavity trees. The land cover throughout the bonneted

220 bat's range includes developed and undeveloped lands, and Bailey et al.'s (2017b) range wide
221 analysis did not find bonneted bats preferentially using pinelands. We believe our analysis at
222 APAFR showed bonneted bat occupancy closely associated with flatwood pinelands because
223 bonneted bats roost in tree cavities of mature pines, and our analysis did not include developed
224 areas that were a minor land cover at APAFR. Although we did not include developed areas as a
225 habitat type for this study, there were multiple ARUs near developed areas that likely would
226 have detected bonneted bats if they occurred in developed areas. We would like to include a
227 covariate of distance to developed lands that includes areas off APAFR as a follow-up analysis.

228 The inclusion of prairie habitats for bonneted bat occupancy may allude to bonneted bats'
229 need for open foraging habitat (Voigt and Holderied 2012). Bonneted bats frequently use
230 agricultural lands, likely as feeding habitats, and the most-similar, open habitat types on APAFR
231 are prairie grasslands (Bailey et al. 2017b). The nearest agricultural lands at APAFR are orange
232 groves, and distance to nearest orange grove was an informative, albeit poorly estimated,
233 covariate for estimating bonneted bats occupancy. We believe that prairie habitats and nearby
234 orange groves provide necessary, proximate insect-rich feeding sites (Simanton 1960; Swengel
235 2001). The loss of southern longleaf and slash pine forests, and suppression of low-intensity,
236 short-return fire intervals that limited understory development in most remaining forests, has
237 greatly reduced the availability of open prairies in mature forests (Croker 1987). Broad swaths of
238 recently-burned forest, with limited vegetative clutter, may have been the primary hunting
239 grounds of bonneted bats. Agricultural lands may now act as surrogate hunting resources for
240 molossid bats (Cleveland et al. 2006; Noer et al. 2012), which may explain their inclusion in
241 bonneted bat occupancy analyses. However, agricultural lands may not provide lepidopteran
242 prey that molossid bats prefer (Krauel et al. 2018). These habitats may not include as much moth

243 diversity and biomass as natural, fire-maintained southern pine grasslands (Armitage and Ober
244 2012).

245 Pine-dominated landscapes cover a large proportion (84%) of APAFR, but only a small
246 fraction (~ 13%) of these pinelands have the mature longleaf and slash pines that would provide
247 roosting habitat for bonneted bats. Currently, there is one active bonneted bat roost, one inactive
248 roost, and three roosts that are no longer viable. All roosts are in woodpecker cavities in mature
249 longleaf pines, except a temporary roost that was under loose bark. The inviable roosts are in
250 trees damaged by hurricanes, in cavities degraded by woodpecker activity, or in use by other
251 species, such as big brown bats *Eptesicus fuscus* or Brazilian free-tailed bats. Efforts to increase
252 bonneted bat habitat at APAFR will require expanding the availability of mature pinelands for
253 the woodpeckers that create bonneted bat roosts. Actively managing forests with historic burning
254 regimes will increase forest types and structure that supports habitat for bonneted bats and other
255 rare species (Van Lear et al. 2005; Braun de Torrez et al. 2018a; 2018b). Restoration of longleaf
256 pine forests at APAFR using natural fire regimes and appropriate silviculture practices
257 (Brockway et al. 2005) will increase habitat suitability for bonneted bats by increasing expanses
258 of mature pines for roost availability, open space for hunting prey, and the insect communities
259 they feed on (Braun de Torrez 2018b).

260 Interestingly, all active and inactive bonneted bat roosts are only along the northern edge of
261 APAFR where mature pinelands are most abundant. Bonneted bat distribution models suggest
262 additional habitat exists in the southern sections of APAFR and Polk County (Bailey et al.
263 2017b). Expanding mature longleaf forests southward would increase the availability and
264 viability of habitat for bonneted bats, and RCW that create cavity roosts. Contiguous forests
265 would increase the success of recovery efforts by connecting bonneted bat habitat and

266 populations, and increasing security from stochastic weather events, such as the increasing
267 frequency and severity of hurricanes (Zampieri et al. 2020). Besides creating forests more
268 attractive to the woodpeckers that create natural cavities, artificial bat boxes have proven to be
269 viable bonneted bat roosting alternates (Bailey et al. 2017a). However, even in areas where bat
270 boxes have been successful, there is an abundance of flatwood forests similar to the mature pine
271 stands of APAFR (FWC 2014). Installation of bat boxes in mature pine stands may provide
272 roosting alternatives for bonneted bats when RCWs are unavailable to build natural roosts.

273 This study is the first to assess bonneted bat habitat use at APAFR; however, because we did
274 not deploy ARUs in all available habitats we cannot assess habitat use in unsampled land cover
275 types (developed areas). Although bonneted bats tend to avoid developed areas (Bailey et al.
276 2017b), future effort should include sampling in these habitats to understand use patterns at
277 APAFR. As further acoustic sampling refines bonneted bat habitat use at APAFR, sampling tools
278 that optimize detection of rare or clustered species can increase efficiency of future sampling
279 design (Brown et al. 2013).

280 Supplemental Material

281 Data S1. Florida bonneted bat *Eumops floridanus* (EUFL) ultrasonic acoustic data from 2018 –
282 2020 from Avon Park Air Force Range, Polk and Highland counties, Florida. The first column
283 represents the location label, and each subsequent column is the detection (1) or failure to detect
284 (0) nightly EUFL acoustic file detection history. Data file (.xlsx)

285
286 Data S2. Site-specific landscape and vegetation covariates from 2018 – 2020 used for modeling
287 Florida bonneted bat *Eumops floridanus* occupancy at Avon Air Force Base, Polk and Highland
288 counties, Florida. The first column represents the location of acoustic recording, and each

289 column represents the landscape- or vegetation-specific attributes of that site. There is a separate
290 tab in the worksheet to define the covariates and their abbreviations. Data file (.xlsx)

291

292 Reference S1. Brockway DG, Outcalt KW, Tomczak DJ, Johnson EE. 2005. Restoration of
293 longleaf pine ecosystems. U.S. Forest Service General Technical Report SRS-83. U.S.
294 Department of Agriculture, Forest Service, Southern Research Station. Asheville, North
295 Carolina. 34 pp.

296 Available: <https://www.srs.fs.usda.gov/pubs/20672>

297

298 Reference S2. Croker TC. 1987. Longleaf pine: a history of man and a forest. United States
299 Department of Agriculture, Forest Service, Forestry Report R8-FR7. U.S. Forest Service,
300 Atlanta, Georgia.

301 Available: <https://www.fs.usda.gov/research/treesearch/59547>

302

303 Reference S3. Florida Fish and Wildlife Conservation Commission (FWC). 2003. A conceptual
304 management plan for Fred C. Babcock – Cecil M. Webb Wildlife Management Area. Florida
305 Fish and Wildlife Conservation Commission, Tallahassee.

306 Available: <https://chnep.wateratlas.usf.edu/upload/documents/cmp-babcock-webb.pdf>

307

308 Reference S4. Plowright A. 2018. ForestTools: Analyzing Remotely Sensed Forest Data. R
309 package version 0.2.0.

310 Available: <https://CRAN.R-project.org/package=ForestTools>.

311

312 Reference S5. Reichert B, Lausen C, Loeb S, Weller T, Allen R, Britzke E, Hohoff T, Siemers J,
313 Burkholder B, Herzog C., Verant M. 2018. A guide to processing bat acoustic data for the North
314 American Bat Monitoring Program (NABat): U.S. Geological Survey Open-File Report 2018–
315 1068, 33 pp.

316 Available: <https://pubs.er.usgs.gov/publication/ofr20181068>

317

318 Reference S6. Su Y, Yajima M. 2021. R2jags: using R to run ‘JAGS’ R package. Version 0.7-1.

319 Available: <https://cran.r-project.org/web/packages/R2jags/index.html>

320

321 Reference S7. [USFWS] U.S. Fish and Wildlife Service. 2013. Endangered species status for the
322 Florida bonneted bat. Federal Register 78:61003-61043.

323 Available: <https://www.federalregister.gov/documents/2013/10/02/2013->

324 [23401/endangered-and-threatened-wildlife-and-plants-endangered-species-status-for-the-florida-](https://www.federalregister.gov/documents/2013/10/02/2013-23401/endangered-and-threatened-wildlife-and-plants-endangered-species-status-for-the-florida-bonneted-bat)
325 [bonneted-bat.](https://www.federalregister.gov/documents/2013/10/02/2013-23401/endangered-and-threatened-wildlife-and-plants-endangered-species-status-for-the-florida-bonneted-bat)

326

327 **Acknowledgments**

328 We are grateful to Emily Angell, Aline Morrow and Katrina Pawlaczyk for their invaluable help
329 in deploying the 500+ bat monitors across APAFR, and Brendan Myers and Troy Hershberger
330 for their role in helping to fund and initiate the acoustic monitoring of bats at APAFR. This
331 research was funded through Fiscal Year 2018 Air Force project AFCW50979318 award to the
332 US Fish and Wildlife Service.

333 Any use of trade, product, website, or firm names in this publication is for descriptive purposes
334 only and does not imply endorsement by the U.S. Government.

References

- 335
336 Agranat I. 2013. Bat species identification from zero crossing and full spectrum echolocation
337 calls using HMMs, Fisher scores, unsupervised clustering, and balanced pairwise classifiers.
338 *Journal of the Acoustical Society of America* 133:331. doi.org/10.1121/1.4799403 (March
339 2023).
- 340 Angell EN, Thompson G. 2015. Second record of natural Florida bonneted bat (*Eumops*
341 *floridanus*) roost in a longleaf pine (*Pinus palustris*). *Florida Field Naturalist* 43:185-188.
- 342 Armitage DW, Ober HK. 2012. The effects of prescribed fire on bat communities in the longleaf
343 pine sandhills ecosystem. *Journal of Mammalogy* 93:102-114. doi.org/10.1644/11-MAMM-
344 A-169.1 (March 2023).
- 345 Austin JD, Gore J, Hargrove J, Braun de Torrez E, Carneiro C, Ridgely F, Wisely S. 2022.
346 Strong population genetic structure and cryptic diversity in the Florida bonneted bat (*Eumops*
347 *floridanus*). *Conservation Genetics* 23:495-512. doi.org/10.1007/s10592-022-01432-y
348 (March 2023)
- 349 Bailey AM, McCleery RA, Ober HK, Pine WE. 2017a. First demographic estimates for
350 endangered Florida bonneted bats suggest year-round recruitment and low apparent survival.
351 *Journal of Mammalogy* 98: 551-559. doi.org/10.1093/jmammal/gyw198 (March 2023)
- 352 Bailey AM, Ober HK, Sovie A, McCleery RA. 2017b. Impact of land use and climate on the
353 distribution of the endangered Florida bonneted bat. *Journal of Mammalogy* 98:1586–1593.
354 doi.org/10.1093/jmammal/gyx117 (March 2023).
- 355 Belwood, J.J. 1992. Florida mastiff bat, *Eumops glaucinus floridanus*. Pages 216 – 232 in
356 Humphrey SR, editor. *Rare and endangered biota of Florida*. Volume I. Mammals.
357 Gainesville, Florida: University Press of Florida.

- 358 Beucher S, Meyer F. 1993. *The morphological approach to segmentation: the watershed*
359 *transformation*. Mathematical Morphology in Image Processing. Baton Rouge, Louisiana:
360 CRC Press.
- 361 Braun de Torrez EC, Gore JA, Ober HK. 2020. Evidence of resource-defense polygyny in an
362 endangered subtropical bat, *Eumops floridanus*. *Global Ecology and Conservation*
363 24:e01289. doi.org/10.1016/j.gecco.2020.e01289 (March 2023).
- 364 Braun de Torrez EC, Ober HK, McCleery RA. 2016. Use of a multi-tactic approach to locate an
365 endangered Florida bonneted bat roost. *Southeastern Naturalist* 15:235–242.
366 doi.org/10.1656/058.015.0204 (March 2023).
- 367 Braun de Torrez EC, Ober HK, McCleery RA. 2018a. Activity of an endangered bat increases
368 immediately following prescribed fire. *Journal of Wildlife Management* 82:1115–1123.
369 doi.org/10.1002/jwmg.21481 (March 2023).
- 370 Braun de Torrez EC, Ober HK, McCleery RA. 2018b. Restoring historical fire regimes increases
371 activity of endangered bats. *Fire Ecology* 14:9. doi.org/10.1186/s42408-018-0006-8 (March
372 2023).
- 373 Brockway DG, Outcalt KW, Tomczak DJ, Johnson EE. 2005. *Restoration of longleaf pine*
374 *ecosystems*. U.S. Forest Service General Technical Report SRS-83. U.S. Department of
375 Agriculture, Forest Service, Southern Research Station. Asheville, North Carolina. 34 pp.
376 (see *Supplemental Material*, Reference S1).
- 377 Brown JA, Salehi MM, Moradi M, Panahbehagh B, Smith DR. 2013. Adaptive survey designs
378 for sampling rare and clustered populations. *Mathematics and Computers in Simulation*
379 93:108-116. doi.org/10.1016/j.matcom.2012.09.008 (March 2023).

- 380 Cleveland CJ, Betke M, Federico P, Frank JD, Hallam TG, Horn J, Lopez, Jr., JD, McCracken
381 GF, Medellín, Moreno-Valdez A, Sansone CG, Westbrook JK, Kunz TH. 2006. Economic
382 value of the pest control services provided by Brazilian free-tailed bats in south-central
383 Texas. *Frontiers in Ecology and the Environment* 4:238-243. doi.org/10.1890/1540-
384 9295(2006)004[0238:EVOTPC]2.0.CO;2 (March 2023).
- 385 Corcoran AJ, Conner WE. 2014. Bats jamming bats: food competition through sonar
386 interference. *Science* 346:745–747. DOI: 10.1126/science.125951 (March 2023).
- 387 Croker TC. 1987. *Longleaf pine: a history of man and a forest*. United States Department of
388 Agriculture, Forest Service, Forestry Report R8-FR7. U.S. Forest Service, Atlanta, Georgia.
389 (see *Supplemental Material*, Reference S2).
- 390 Duever MJ, Meeder JF, Meeder LC, McCollom JM. 1994. The climate of South Florida and its
391 role in shaping the Everglades ecosystem. Pages 225-248 in Davis SM, Ogden JC, editors.
392 *Everglades: the ecosystem and its restoration*. Delray Beach, Florida: St. Lucie Press.
- 393 [ESA] US Endangered Species Act of 1973, as amended, Pub. L. No. 93-205, 87 Stat. 884 (Dec.
394 28, 1973). Available at: <https://www.fws.gov/endangered/esa-library/pdf/ESAall.pdf>
- 395 Florida Fish and Wildlife Conservation Commission (FWC). 2003. *A conceptual management*
396 *plan for Fred C. Babcock – Cecil M. Webb Wildlife Management Area*. Florida Fish and
397 Wildlife Conservation Commission, Tallahassee. (see *Supplemental Material*, Reference S3).
- 398 Gore JA, Robson MS, Zambrano R, Douglass NJ. 2015. Roosting sites of a Florida bonneted bat
399 (*Eumops floridanus*). *Florida Field Naturalist* 43:179-184.
- 400 Hooten MB, Hobbs NT. 2015. A guide to Bayesian model selection for ecologists. *Ecological*
401 *Monographs* 85:3-28. doi.org/10.1890/14-0661.1 (March 2023).

- 402 Jones TK, Wohlgemuth MJ, Conner WE. 2018. Active acoustic interference elicits echolocation
403 changes in heterospecific bats. *Journal of Experimental Biology* 221:jeb176511.
404 doi.org/10.1242/jeb.176511 (March 2023).
- 405 Kery M, Royle JA. 2015. *Applied hierarchical modeling in ecology: analysis of distribution,*
406 *abundance, and species richness in R and BUGS.* Academic Press, London, United
407 Kingdom.
- 408 Krauel JJ, Ratcliffe JM, Westbrook JK, McCracken GF. 2018. Brazilian free-tailed bats
409 (*Tadarida brasiliensis*) adjust foraging behavior in response to migratory moths. *Canadian*
410 *Journal of Zoology* 96:513-520. doi.org/10.1139/cjz-2017-0284 (March 2023).
- 411 Kuo L, Mallick B. 1998. Variable selection for regression models. *The Indian Journal of*
412 *Statistics: Special Issue on Bayesian Analysis* 60:65-81.
413 <https://www.jstor.org/stable/25053023> (March 2023)
- 414 Morin DJ, Yackulic CB, Diffendorfer JE, Lesmeister DB, Nielsen CK, Reid J, Schauber EM.
415 2020. Is your ad hoc model selection strategy affecting your multimodel inference?
416 *Ecosphere* 11:e020997. doi.org/10.1002/ecs2.2997 (March 2023).
- 417 Noer CL, Dabelsteen T, Bohmann K, Monadjem A. 2012. Molossid bats in an African agro-
418 ecosystem select sugarcane fields as foraging habitat. *African Zoology* 47:1-11.
419 doi.org/10.1080/15627020.2012.11407517 (March 2023).
- 420 Ntzoufras, I. 2009. *Bayesian modeling using WinBUGS.* Wiley, Hoboken, New Jersey.
- 421 Plowright A. 2018. ForestTools: analyzing remotely sensed forest data. R package version 0.2.0.
422 (see *Supplemental Material*, Reference S4).
- 423 Popescu SC, Wynne RH. 2004. Seeing the trees in the forest. *Photogrammetric Engineering &*
424 *Remote Sensing* 70:589-604. doi.org/10.14358/PERS.70.5.589 (March 2023).

- 425 R Core Team. 2018. *R: A language and environment for statistical computing*. R Foundation for
426 Statistical Computing, Vienna, Austria.
- 427 Reichert B, Lausen C, Loeb S, Weller T, Allen R, Britzke E, Hohoff T, Siemers J, Burkholder B,
428 Herzog C., Verant M. 2018. *A guide to processing bat acoustic data for the North American
429 Bat Monitoring Program (NABat): U.S. Geological Survey Open-File Report 2018–1068*, 33
430 pp. (see *Supplemental Material*, Reference S5).
- 431 Royle JA, Dorazio RM. 2008. *Hierarchical modeling and inference in ecology*. Elsevier,
432 London, United Kingdom.
- 433 Simanton WA. 1960. Seasonal populations of citrus insects and mites in commercial groves. *The
434 Florida Entomologist* 43:49-57. doi.org/10.2307/3492379 (March 2023).
- 435 Slocum MG, Platt WJ, Beckage B, Orzell SL, Taylor W. 2010. Accurate quantification of
436 seasonal rainfall and associated climate–wildfire relationships. *Journal of Applied
437 Meteorology and Climatology* 49:2559–2573. doi.org/10.1175/2010JAMC2532.1 (March
438 2023).
- 439 Smith DR, Allan NL, McGowan CP, Szymanski JA, Oetker SR, Bell HM. 2018. Development of
440 a Species Status Assessment process for decisions under the U.S. Endangered Species Act.
441 *Journal of Fish and Wildlife Management* 9:302-320. https://doi.org/10.3996/052017-JFWM-
442 041 (March 2023)
- 443 Su Y, Yajima M. 2021. R2jags: using R to run ‘JAGS’ R package. Version 0.7-1. (see
444 *Supplemental Material*, Reference S6).
- 445 Swengel AB. 2001. A literature review of insect responses to fire, compared to other
446 conservation managements of open habitat. *Biodiversity and Conservation* 10:1141-1169.
447 doi.org/10.1023/A:1016683807033 (March 2023).

- 448 [USFWS] U.S. Fish and Wildlife Service. 2013. Endangered species status for the Florida
449 bonneted bat. *Federal Register* 78:61003-61043. (see *Supplemental Material*, Reference S7).
- 450 Van Lear DH, Carroll WD, Kapeluck PR, Johnson R. 2005. History and restoration of the
451 longleaf pine-grassland ecosystems: implications for species at risk. *Forest Ecology and*
452 *Management* 211:150-165. doi.org/10.1016/j.foreco.2005.02.014 (March 2023).
- 453 Voigt CC, Holdereid MW. 2012. High manoeuvring costs force narrow-winged molossid bats to
454 forage in open space. *Journal of Comparative Physiology B* 182:415-424.
455 doi.org/10.1007/s00360-011-0627-6 (March 2023).
- 456 Walker KJ. 2000. Historical ecology of the southeastern longleaf and slash pine flatwoods: a
457 southwest Florida perspective. *Journal of Ethnobiology* 20:269-299.
458 <https://ethnobiology.org/sites/default/files/pdfs/JoE/20-2/Walker.pdf> (March 2023).
- 459 Webb, EN, Ober HK, Braun de Torrez EC, Gore JA, Zambrano R. 2021. Urban roosts: use of
460 buildings by Florida bonneted bats. *Urban Naturalist* 42:1-11. [https://eaglehill.us/urna-pdfs-](https://eaglehill.us/urna-pdfs-regular/urna-042-Ober.pdf)
461 [regular/urna-042-Ober.pdf](https://eaglehill.us/urna-pdfs-regular/urna-042-Ober.pdf) (March 2023).
- 462 Zampieri NE, Pau S, Okamoto DK. 2020. The impact of Hurricane Michael on longleaf pine
463 habitats in Florida. *Scientific Reports* 10:8483. doi.org/10.1038/s41598-020-65436-9 (March
464 2023).

465

466 Figure 1. Photographs of the Florida bonneted bat *Eumops floridanus* and bonneted bat habitat
467 from southern Florida, including (A), a Florida bonneted from Fred C. Babcock/Cecil M. Webb
468 Wildlife Management Area, Florida, 2018, (B), typical longleaf pine *Pinus palustris* forest at
469 Avon Park Air Force Range (APAFR), Florida, including, and (C) biologists looking into a
470 bonneted bat roost, APAFR, 2019. Photographs taken by RAS, KAP.

471

472 Figure 2. Map of Avon Park Air Force Range location in Florida and the landcover types and
473 autonomous recording unit (ARU) locations on Avon Park Air Force Range, 2018 – 2020.

474

475 Figure 3. Florida bonneted bat *Eumops floridanus* occupancy and distance to red-cockaded
476 woodpecker *Picoides borealis* colony at Avon Park Air Force Range, Florida, 2018-2019.

477

478

479 Table 1. Covariates and their predicted impact on occupancy and detectability of Florida
480 bonneted bat *Eumops floridanus* at Avon Park Air Force Range, Avon Park, Florida, 2018 -
481 2019.

482


483 Table 2. Covariates with $\geq 50\%$ inclusion in variable selection runs and estimated effect size (β)
484 and 95% credible interval (CI) for occupancy and detectability of Florida bonneted bats *Eumops*
485 *floridanus* at Avon Park Air Force Range, Florida, 2018 - 2019.

486

<Fig. 1>



<Fig. 2>



<Fig. 3>




Table 1. Covariates and their predicted impact on occupancy and detectability of F
Air Force Range, Avon Park, Florida, 2018 - 2019.

Covariate	How applied
ARU Nightly Temperature (mean, maximum, minimum)	Daily
Scotophase	Daily
Nightly Temperature (mean, maximum, minimum)	Daily
Nightly Precipitation	Daily
Nightly wind (mean, maximum, minimum)	Daily
<i>Tadarida brasiliensis</i> activity	Daily
Acoustic noise	Daily
Distance from nearest <i>Picoides borealis</i> cluster	Site
Distance from nearest <i>Citrus sinensis</i> grove	Site
Area of nearest <i>Citrus sinensis</i> grove	Site
Distance from nearest wetland	Site
Area of nearest wetland	Site
Habitat type	Site
Distance from nearest tree within 100 m	Site
Height of nearest tree within 100 m	Site
Canopy radius (m) of nearest tree within 100 m	Site
Canopy cover within 100 m	Site
Percent canopy cover within 100 m	Site
Crown area (m ²) within 100 m (mean, maximum, minimum)	Site
Height of trees within 100 km (mean, maximum, minimum)	Site
Canopy radius (m) of trees within 100 km (mean, maximum, minimum)	Site
Days since last forest fire	Site

Florida bonneted bat *Eumops floridanus* at Avon Park


Predicted impact	
Occupancy	Detectability
-	increase
-	decrease
-	increase
-	decrease
-	decrease
-	decrease
-	decrease
decrease	-
decrease	-
increase	-
increase	increase
increase	increase
decrease/increase	decrease/increase
increase	-
increase	-
decrease	-
decrease	-
decrease	-
decrease	-
decrease	-
decrease	-
decrease	-

Table 2. Covariates with $\geq 50\%$ inclusion in variable selection runs and estimated effect size (β) and 95 (CI) for occupancy and detectability of Florida bonneted bats *Eumops floridanus* at Avon Park Air For 2018 - 2019.



Detectability	Percent inclusion	β (95% CI)
Habitat - Flatwoods	100	-2.52 (-2.74, -2.32)
Habitat - Grass/Prairie	50	-2.67 (-3.11, -2.28)
Habitat - Oak	100	-4.19 (-5.54, -2.84)
Habitat - Pine Plantation	100	-2.65 (-3.04, -2.28)
Habitat - Scrub	100	-2.39 (-2.89, -1.93)
Habitat - Swamp/Marsh	100	-2.33 (-3.24, -1.58)
Julian Date	100	0.41 (0.27, 0.55)
<hr/>		
Occupancy		
Habitat - Flatwoods	100	-1.25 (-1.72, -0.72)
Habitat - Grass/Prairie	57	-0.55 (-8.80, 8.82)
Habitat - Pine Plantation	53	-0.47 (-8.87, 8.92)
Area of nearest wetland	50	-0.26 (-8.80, 9.00)
Distance to nearest <i>Citrus sinensis</i> grove	72	-0.43 (-8.17, 8.15)
Distance to nearest <i>Picoides borealis</i> colony	100	-1.19 (-1.70, -0.72)


% credible interval
rice Range, Florida,

<u>parameter (95% CI)</u>
0.08 (0.06, 0.09)
0.07 (0.04, 0.09)
0.02 (0.00, 0.06)
0.07 (0.05, 0.09)
0.09 (0.05, 0.13)
0.10 (0.04, 0.17)
0.60 (0.57, 0.63)
0.23 (0.15, 0.33)
0.38 (0.00, 1.00)
0.39 (0.00, 1.00)
0.44 (0.00, 1.00)
0.40 (0.00, 1.00)
0.24 (0.15, 0.33)






Click here to access/download
Supplemental Material
S1.xlsx





Click here to access/download
Supplemental Material
S2.xlsx






Click here to access/download
Supplemental Material
Brockway et al. 2005 restoration of longleaf pine
ecosystems.pdf





Click here to access/download
Supplemental Material
FWC 2003 Conceptual management plan BWMA.pdf





Click here to access/download
Supplemental Material
Reichert et al 2018.pdf



Click here to access/download
Supplemental Material
R2jags.pdf



Click here to access/download

Supplemental Material

USFWS ES status Florida bonneted bat.pdf

